HYDROBIOLOGIA
Volume 449, 15 April 2001

Theme
Advances in Decapod Crustacean Research
Proceedings of the 7th Colloquium Crustacea Decapoda Mediterranea, 6–9 September 1999

Guest Editors
José Paula, Augusto Flores & Charles Fransen

Preface ix

1. SYSTEMATICS, PHYLOGENY AND BIOGEOGRAPHY

Phylogeny of decapods: moving towards a consensus
F.R. Schram 1–20

Notes on the position of the true freshwater crabs within the Brachyrhynchan Eubrachyura (Crustacea: Decapoda: Brachyura)
R.v. Sternberg, N. Cumberlidge 21–39

Molecular phylogeny of the crab genus *Brachynotus* (Brachyura: Varunidae) based on the 16S rRNA gene
C.D. Schubart, J.A. Cuesta, A. Rodríguez 41–46

***Austinogebia*, a new genus in the Upogebidae and rediagnosis of its close relative, *Gebiacantha* Ngoc-Ho, 1989 (Crustacea: Decapoda: Thalassinidea)**
N. Ngoc-Ho 47–58

Recent samples of mainly rare decapod Crustacea taken from the deep-sea floor of the southern West Europe Basin
L. Tiefenbacher 59–70

Data on the family Pandalidae around the Canary Islands, with first record of *Plesionika antigai* (Caridea)
J.A. González, J.A. Quiles, V.M. Tuset, M.M. García-Díaz, J.I. Santana 71–76

Crustacea Decapoda of Paripe River Estuary, Pernambuco, Brazil
M.A. Coelho dos Santos, P.A. Coelho 77–79

Intertidal habitats and decapod faunal assemblages (Crustacea: Decapoda) of Socotra Island, Republic of Yemen
N. Simões, M. Apel, D.A. Jones 81–97

The crab species found on the coasts of Gökçeada (Imbroz) Island in the Aegean Sea
H. Balkıs, N. Balkıs, S. Altınsaçlı 99–103

Evidence of paraphyly in the neotropical Porcellanid genus *Neopisosoma* (Crustacea: Anomura: Porcellanidae) based on molecular characters
B. Werding, A. Hiller, B. Misof 105–110

2. GROWTH, MORPHOLOGY AND DEVELOPMENT

Growth in Crustacea – twenty years on
R.G. Hartnoll 111–122

Sex-related variability of rostrum morphometry of *Aristeus antennatus* (Decapoda: Aristeidae) from the Ionian Sea (Eastern Mediterranean, Greece)
K. Kapiris, M. Thessalou-Legaki 123–130

Influence of diet on sex differentiation of *Hippolyte inermis* Leach (Decapoda: Natantia) in the field
V. Zupo 131–140

Combined effects of temperature and salinity on the larval development of the estuarine mud prawn *Upogebia africana* (Crustacea, Thalassinidea)
J. Paula, R. Nogueira Mendes, S. Paci, P. McLaughlin, F. Gherardi, W. Emmerson 141–148

Mitotic and meiotic chromosomes of the American lobster *Homarus americanus* (Nephropidae, Decapoda)
E. Coluccia, A. Cau, R. Cannas, A. Milia, S. Salvadori, A.M. Deiana 149–152

3. ECOLOGY AND BEHAVIOUR

Larval abundance and recruitment of *Carcinus maenas* L. close to its southern geographic limit: a case of match and mismatch
M. Sprung 153–158

Choice of prey size and species in *Carcinus maenas* (L.) feeding on four bivalves of contrasting shell morphology
M. Mascaró, R. Seed 159–170

Intertidal distribution and species composition of brachyuran crabs at two rocky shores in central Portugal
A.A.V. Flores, J. Paula 171–177

Morphometric comparison between Mediterranean and Atlantic populations of *Pontophilus norvegicus* (Decapoda, Crangonidae)
S. De Grave, D. Diaz 179–186

Notes on the distribution and biology of the deep-sea crab *Bathynectes maravigna* (Brachyura: Portunidae) in the Mediterranean Sea
P. Abelló, N. Ungaro, C.-Y. Politou, P. Torres, E. Roman, P. Rinelli, P. Maiorano, G. Norrito 187–192

Notes on the biology of *Cancer bellianus* (Brachyura, Cancridae) around the Canary Islands
J.A. Quiles, V. Rico, V.M. Tuset, J.I. Santana, J.A. González 193–199

Comparative suitability of binocular observation, burrow counting and excavation for the quantification of the mangrove fiddler crab *Uca annulipes* (H. Milne Edwards)
M.W. Skov, R.G. Hartnoll 201–212

A comparison of alternative methods for estimating population density of the fiddler crab *Uca annulipes* at Saco Mangrove, Inhaca Island (Mozambique)
A. Macia, I. Quincardete, J. Paula 213–219

Aspects of the population dynamics of *Neosarmatium meinerti* at Mgazana, a warm temperate mangrove swamp in the East Cape, South Africa, investigated using an indirect method
W.D. Emmerson 221–229

Seasonal abundance and recruitment in an estuarine population of mud crabs, *Scylla paramamosain*, in the Mekong Delta, Vietnam
L. Le Vay, V.N. Ut, D.A. Jones 231–239

Major claws make male fiddler crabs more conspicuous to visual predators: a test using human observers
J.M. Jordão, R.F. Oliveira 241–247

Feeding activity of *Callinectes ornatus* Ordway, 1863 and *Callinectes danae* Smith, 1869 (Crustacea, Brachyura, Portunidae) in Ubatuba, SP, Brazil
A.L.D. Reigada, M.L. Negreiros-Fransozo 249–252

4. REPRODUCTION

A histochemical and ultrastructural study of oogenesis in *Aristaeomorpha foliacea* (Risso, 1827)
S. Desantis, M. Labate, P. Maiorano, A. Tursi, G.M. Labate, M. Ciccarelli 253–259

Prevalence of bacteria in the spermathecae of female snow crab, *Chionoecetes opilio* (Brachyura: Majidae)
K. Benhalima, M. Moriyasu 261–266

5. FISHERIES AND CULTURE

Discards of the Algarve (southern Portugal) crustacean trawl fishery
P. Monteiro, A. Araújo, K. Erzini, M. Castro 267–277

Effect of codend mesh size on the performance of the deep-water bottom trawl used in the red shrimp fishery in the Strait of Sicily (Mediterranean Sea)
S. Ragonese, M. Zagra, L. Di Stefano, M.L. Bianchini 279–291

Protein requirement of the prawn *Marsupenaeus japonicus* estimated by a factorial method
S. Teshima, S. Koshio, M. Ishikawa, A. Kanazawa 293–300

The effect of fixatives in the quantification of morphological lipofuscin as an age index in crustaceans
P. Encarnação, M. Castro 301–305